

# PROXIMATE, MINERAL AND PHYTOCHEMICAL ANALYSIS OF SOME MEDICINAL PLANTS COLLECTED FROM ORATHUR VILLAGE, THIRUPORUR TALUK KANCHEEPURAM DISTRICT TAMILNADU, INDIA

Alagbe Olujimi John

Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India.

Corresponding Email: dralagbe@outlook.com

**Abstract:** The use of medicinal plants has a long tradition as part of the human diet and has therapeutic agents. That is why they are well accepted by consumers and are generally considered to be environmentally friendly, efficient and generally considered to be safe alternative to antibiotics due to the presence of phytochemicals found in the leaves, stem bark, roots, flowers amongst others. Medicinal plant also possesses nutritional properties and play multiple pharmacological roles such as anti-inflammatory, antioxidant, anti-cancer, antimicrobial, anti-viral, anti-fungal, anti-helminthic, hepato-protective activity amongst others. This study was aimed at accessing the proximate, mineral and phytochemical composition of *Tephrosia purpurea*, *Adhatoda vasica*, *Crataeva nurvala*, *Bauhinia racemose*, *Holarrhena antidysenterica* and *Aegle marmelos*. Result on proximate analysis showed that moisture content varied from 13.57 – 16.02 %, dry matter (83.98 – 86.43 %), crude protein (5.06 – 7.12 %), ash (9.90 – 12.04 %), crude fibre (12.74 – 14.32 %), crude fat (0.10 – 0.22 %) and carbohydrates (40.30 – 50.17 %). Concentrations of calcium, phosphorus, potassium, magnesium, zinc, manganese, sodium and copper ranged from 120 – 148.0 mg/100g, 180.0 – 300 mg/100g, 66.8 – 94.2 mg/100g, 41.8 – 69.8 mg/100g, 27.9 – 44.6 mg/100g, 119 – 133 mg/100g and 8.21 – 13.1 mg/100g respectively. Quantitative evaluation of tannins showed that values varied from 39.00 – 46.50 mg/g, alkaloids (26.35 – 35.09 mg/g), saponins (20.05 – 30.40 mg/g), flavonoids (66.10 – 86.11 mg/g), terpenoids (40.71 – 45.88 mg/g) and phenols (106.8 – 125.4 mg/g). In conclusion, it was discovered that it contained some essential nutrients in significant quantities. However, values obtained varied due to age of plant, specie, processing method amongst others. The presence of phytochemicals in these plant suggests that it has pharmacological properties. Accessing the constituent's in this plants could lead to a novel drug discovery and curb the increasing cases in antimicrobial resistance.

**Keywords:** Antimicrobials; Plants; Therapeutic; Food safety; Phytochemicals

## 1 INTRODUCTION

Growing antibiotic resistance is leading to a continuous need for discovering new drugs and alternative treatment against diseases (Vera, 2022)[1]. Antimicrobial resistance poses a serious hazard to the populace making it difficult for diseases to be treated and also increasing mortality and morbidity (Vera, 2022[1]; Ojediran et al., 2024[2]). In 2009, the World Health Organization named antimicrobial resistance as one of the three greatest threats to human health and in 2011 the focus of the World Health day was “Combating antimicrobial resistance”.

One of the alternatives that can be utilized for combating antimicrobial resistance is traditional herbal remedies or the use of medicinal plants (Inge and Rainer, 2020)[4]. Medicinal plants possess nutritional and pharmacological properties and have been used for the treatment of various ailments due to the presence of phyto-constituents or phytochemicals (alkaloids, tannins, flavonoids, terpenoids, phenols, saponins and so on) (Ojediran et al., 2024[2]; Singh et al., 2022[5]). According to John (2024)[6]; Inge and Raine (2020)[4], phytochemicals are eco-friendly, effective, safe and has no withdrawal period. They are found in many parts of plants including, root, stem bark, leaves, flowers, buds, amongst others and capable of performing multiple therapeutic activities (anti-inflammatory, antifungal, anti-atherosclerotic, immuno-stimulatory, antioxidant, hypolipidemic, cytotoxic, anti-parasitic and anti-thrombotic) in human beings and livestock (Alagbe, 2022[34]; Musa et al., 2020)[8].

*In vitro* studies have revealed that phytochemicals can inhibit the activities of some pathogenic organisms such as, *Escherichia coli*, *Salmonella spp*, *Staphylococcus spp*, *Aspergillus spp*, *Candida albicans*, *Bacillus spp* by perforating the lipid bilayer of these pathogens, increasing the cell wall permeability thus damaging their genetic material and preventing cell replication. Qualitative and quantitative variations in the contents of nutrients and phytochemicals in plant materials can be influenced by age of plant, geographical locations, climate, species, method of processing and storage (Ahmad et al., 2006)[9].

Therefore, this study was carried out to examine the proximate, mineral and phytochemical analysis of some medicinal plants collected from Orathur village, Thiruporur Taluk Kancheepuram district Tamilnadu, India.

## 2 MATERIALS AND METHODS

### 2.1 Experimental Location

This experiment was carried out at the department of biochemistry, Sumitra Research Institute, Gujarat India situated between 28° 18' N and 70° 35' E India in the month of October 2022. Kits/ equipment's used for the study were handled based on the manufacturers recommendation on each package insert.

### 2.2 Collection of Medicinal Plants and Their Processing

Fresh leaves of medicinal plants (*Tephrosia purpurea*, *Adhatoda vasica*, *Crataeva nurvala*, *Bauhinia racemose*, *Holarrhena antidysenterica* and *Aegle marmelos*) were collected from different sites at Orathur village, Thiruporur Taluk Kancheepuram district Tamilnadu, India and sent to the department of Crop protection at Sumitra Research Institute, Gujarat for proper identification by a certified taxonomist and assigned individual voucher numbers: AA/2022DA, AB/2022DA, AC/2022DA, AD/2022DA, AE/2022DA and AF/2022DA respectively. Thereafter, plants were sorted into different groups, washed under running tap water and air dried for 2 weeks until a constant weight was achieved. Dried leaves were grinded into powder with an electric blender and stored in a clean labeled airtight plastic container before it was sent to the laboratory for further evaluation.

### 2.3 Proximate Analysis of Samples

Proximate composition of each medicinal plant were determined using diode array based near infra-red reflectance and trans reflectance analyzer (Model NIRSTM DA 1650). 100 g of each sample is placed in the collector after the machine was calibrated following the manufacturers lay down procedures. To maintain further precision, it was adjusted at an optical bandwidth (8.75 nm), spectral resolution (1.0 nm), absorbance ranges (up to 2 AU), wavelength accuracy (less than 0.05 nm), photometric noise (400 – 700 nm less 50 micro au; 700 – 2500 nm less than 20 micro au) before results on moisture content, crude protein, crude ash, ether extract and carbohydrate was generated via the visual display unit in 60 seconds. Dry matter was computed by subtracting the moisture content in each sample from 100.

### 2.4 Mineral Analysis of Samples

Mineral compositions were analyzed using Atomic absorption spectrophotometer fully automated flame and graphite furnace system (Model AAS – 4000). Each component is maintained at different conditions to ensure precision in results. 100 g of each sample is injected into the machine and adjusted at different specifications; graphite furnace system is transversally heated and maintained at a temperature range of 3000 °C, detection limit (Cd<0.04ng/ml) and sensitivity (50ng/ml absorption >0.40 Abs) while the monochromator was placed at wavelength accuracy, resolution, reproducibility (0.15nm, 0.2nm and <0.05nm), blazed wavelength (250nm), focus (300 nm) and wavelength range (185nm – 1000nm).

### 2.5 Quantitative Evaluation of Phyto-constituents in Samples

Phyto-constituents were analyzed according to the procedures recently published by Alagbe. 100 grams was used for each analysis and each constituent were recorded at different optical density (alkaloids, 500 nm), flavonoids (460 nm), terpenoids (370 nm), alkaloids (550 nm), tannins (480 nm), steroids (250 nm) and phenols (670 nm) using GC-MS 6800 gas chromatography/mass spectrometer. To ensure precision, gas chromatography chamber where the sample is first injected is maintained at an inlet temperature of 450 °C, column temperature (4 – 450 °C, pressure range (0 – 100 psi ± 0.002 psi) and heating rate up to 1201/min while the mass spectrometer unit is adjusted at an ion source temperature (100 – 350 °C), stability (± 0.10 amu/48 hours), mass range (1.5 – 1000 amu), scan rate (up to 10000 amu/sec) and ionization energy (5 eV – 250 eV).

### 2.6 Statistical Analysis

All measurement was carried out in triplicates and results was analyzed using Statistical Package for Social Science (SPSS for Windows (IBM), version 25, Chicago, USA, 2011). Obtained results were expressed as means ± SEM.

## 3 RESULTS AND DISCUSSION

Proximate composition of different medicinal plants collected from Orathur village, India is presented in Table 1. Moisture content obtained in this study varied from 13.57 – 16.02 %, dry matter (83.98 – 86.43 %), crude protein (5.06 – 7.12 %), ash (9.90 – 12.04 %), crude fibre (12.74 – 14.32 %) and carbohydrates (40.30 – 50.17 %). Leaves from *Tephrosia purpurea* had the highest moisture content (16.02 %) followed by *Aegle marmelos* (15.22 %), *Aegle marmelos* (15.10 %), *Holarrhena*

antidysenterica (14.80 %), *Bauhinia racemose* (14.12 %) and *Adhatoda vasica* (13.57 %). The moisture content recorded in this study is similar to those obtained in *Cnidoscopus aurifolia* (12.24 %), *Indigofera tinctoria* (13.60 %), *Daniellia oliveri* stem bark (12.61 %) and *Delonix regia* leaves (14.37 %) reported by Akaninyene and Uwemedimo (2016)[16]; Alagbe et al. (2020)[17]. This suggests that the samples can be stored for a longer period without deteriorating due to microbial growth (Alagbe et al., 2020)[18]. Dry matter values obtained is in agreement with those obtained for *Balanites aegyptiaca* stem bark (88.13 %); *Alchornea cordifolia* stem bark (89.00 %) but lower than those reported for *Chenopodium ambrosioides* (10.60 %), *Morinda lucida* leaves (20.80 %), *Veronia amygdalina* leaves (20.80 %) reported by Aborisade et al. (2017)[10]; Musa et al. (2020)[8]. Crude protein values were highest in *Bauhinia racemose* leaves (7.12 %) while *Tephrosia purpurea* leaves had the lowest concentration (4.93 %). Result obtained for crude protein was similar to those reported for *Balanites aegyptiaca* stem bark (6.60 %) by Musa et al. (2020)[8] but lower than values reported for *Daniellia oliveri* leaves (25.03 %) and *Leptadenia hastata* leaves (17.96 %) recorded by Lawal et al. (2002)[11]. This variation could be due to species, age of plant, geographical location and processing methods employed (Agubosi et al., 2022[12]; Alagbe, 2022[34]). Protein are necessary for growth and the repair of worn-out tissues (Singh et al., 2022[5]). However, values reported was not within the daily recommended intake for birds (8.0 – 12.0 %) (Sales and Janesens, 2003[14]; NRC, 1994)[15]. *Bauhinia racemose* leaves had a higher concentration of 12.04 % followed by *Crataeva nurvala* leaves (11.55 %), *Holarrhena antidysenterica* leaves (11.30 %), *Tephrosia purpurea* (10.69 %), *Adhatoda vasica* (10.41 %) and *Aegle marmelos* leaves (9.90 %). Values obtained were higher than those recorded for *Vitex doniana* leaves (1.04 %), *Cnidoscopus aurifolia* (6.64 %) reported by Lawal et al. (2002)[11]; Akaninyene and Uwemedimo (2016)[16]. This result suggests that *Bauhinia racemose* leaves contains significant quantity of minerals relative to the other plants making them useful in enzymatic activities in the body (Ojediran et al., 2024)[3]. Dietary crude fibre play a significant role in the preventing coronary heart disease and facilitating the digestion of foods (John, 2024)[17]. However, results obtained showed that *Holarrhena antidysenterica* had the highest value of 14.32 % while *Tephrosia purpurea* leaves had the lowest concentration (13.53 %). Results obtained were lower than values recorded for *Crescentia cujete* leaves (19.72 %) by Musbau et al. (2018)[35] but higher than those of *Crataeva nurvala* leaves 6.32 % and *Crataeva nurvala* stem bark (9.72 %) reported by Nahar et al. (2023)[19]. Crude fat values which varied from 0.10 – 0.22 % were higher in *Holarrhena antidysenterica* (0.22 %) and lower in *Bauhinia racemose* leaves (0.10 %). However, values were within the values reported by Nahar et al. (2023)[19] for *Crataeva nurvala* leaves (0.16 %) and *Crataeva nurvala* stem bark (0.12 %). Values for carbohydrates showed that *Bauhinia racemose* had the highest concentration (50.17 %) which is within the daily recommended intake for birds (NRC, 1994)[15] compared to *Holarrhena antidysenterica* which had the lowest value (40.30 %). Values obtained was higher than those recorded for *Crataeva nurvala* leaves (17.68 %) and *Cnidoscopus aurifolia* leaves 2.27 % reported by Nahar et al. (2023)[19]; Akaninyene and Uwemedimo (2016)[16]. Carbohydrates are needed to supply energy for body process and aids the absorption of calcium (Singh et al., 2022[5]).

**Table 1** Proximate composition of different medicinal plants collected from Orathur village, India

Medicinal plants	<sup>1</sup> MC(%)	<sup>2</sup> DM (%)	<sup>3</sup> CP (%)	Ash (%)	<sup>4</sup> CF (%)	<sup>5</sup> CHO (%)	<sup>6</sup> CFF
<i>Tephrosia purpurea</i> leaves	16.02± 0.08	83.98±0.56	4.93±0.71	10.69±1.66	13.53±0.06	42.18±2.33	0.12±0.00
<i>Adhatoda vasica</i> leaves	13.57± 0.45	86.43±0.22	6.99±0.51	10.41±0.97	14.05±0.07	46.22±1.51	0.14±0.01
<i>Crataeva nurvala</i> leaves	15.10± 0.93	84.90±0.18	5.08±0.06	11.55±1.52	13.80±0.43	41.05±1.41	0.18±0.02
<i>Bauhinia racemose</i> leaves	14.12± 0.14	85.88±0.38	7.12±0.19	12.04±1.88	12.74±0.02	50.17±0.97	0.10±0.00
<i>Holarrhena antidysenterica</i> leaves	14.80± 0.64	85.20±0.30	4.95±0.02	11.30±2.84	14.32±0.09	40.30±0.21	0.22±0.06
<i>Aegle marmelos</i> leaves	15.22± 0.42	84.78±0.26	5.06±0.58	9.90±0.73	14.00±0.12	40.96±0.40	0.11±0.00

<sup>1</sup>Moisture content; <sup>2</sup>dry matter; <sup>3</sup>crude protein; <sup>4</sup>crude fibre; <sup>5</sup>carbohydrates; <sup>6</sup>crude fat

Mineral composition of different medicinal plants collected from Orathur village, India expressed in mg/100g is presented in Table 2. *Bauhinia variegata* leaves had the highest concentration of calcium (148 mg/100g), phosphorus (300 mg/100g), potassium (94.2 mg/100g), magnesium (162.0 mg/100g), zinc (69.8 mg/100g), manganese (44.6 mg/100g), sodium (133.0 mg/100g) and copper (13.1 mg/100g) while *Aegle marmelos* had the lowest concentration with calcium (120 mg/100g), phosphorus (180.0 mg/100g), potassium (60.4 mg/100g), magnesium (100.0 mg/100g), zinc (51.7 mg/100g), manganese (31.9 mg/100g), sodium (121.0 mg/100g) and copper (10.2 mg/100g). Values of calcium, phosphorus, potassium,

magnesium, zinc, manganese, sodium and copper in *Tephrosia purpurea*, *Adhatoda vasica*, *Crataeva nurvala*, *Holarrhena antidysenterica* and *Aegle marmelos* leaves varied from 125.0 – 141.0 mg/100g, 180.0 – 291.0 mg/100g, 66.8 – 86.3 mg/100g, 100.0 – 151.0 mg/100g, 41.80 – 61.30 mg/100g, 27.9 – 40.8 mg/100g, 119.0 – 125.0 mg/100g and 8.21 – 12.4 mg/100g respectively. However, all values were within the daily recommended allowance (FDA, 2019[20]; Julian, 2021)[21]. Calcium concentration reported in this study was higher than values recorded for *Hippocratea myriantha* (210 mg/100g,) and *Urera trinervis* leaves (220 mg/100g) reported by Andzouana and Mombouli (2012)[22]. Calcium are needed for the healthy bones in animals. Potassium, magnesium, manganese and zinc values obtained were lower than values reported for *Xylopi aethiopia* 94.11 mg/100g, 0.22 mg/100g, 0.003 mg/100g and 0.020 mg/100g by Abolaji et al. (2007)[23]. Minerals are needed as a co-factor of metabolic enzyme, regulate heart rate, membrane potential, regulate body fluid and maintain pH balance.

**Table 2** Mineral composition of different medicinal plants collected from Orathur village, India (mg/100g)

Medicinal plants	<sup>1</sup> Ca	<sup>2</sup> P	<sup>3</sup> K	<sup>4</sup> Mg	<sup>5</sup> Zn	<sup>6</sup> Mn	<sup>7</sup> Na	<sup>8</sup> Cu
<i>Tephrosia purpurea</i> leaves	125±2.17	233±2.4	82.1±1.8	112±4.9	61.3±0.8	30.2±1.8	123±6.60	11.2±0.01
<i>Adhatoda vasica</i> leaves	140±2.00	291±4.2	78.5±2.0	109±5.6	56.4±0.9	34.5±1.5	125±5.57	10.9±0.00
<i>Crataeva nurvala</i> leaves	137±1.82	262±3.7	66.8±2.7	151±5.3	42.1±0.3	40.8±1.0	119±3.10	12.4±0.04
<i>Bauhinia variegata</i> leaves	148±1.61	300±2.5	94.2±2.5	162±4.1	69.8±0.5	44.6±0.9	133±2.84	13.1±0.01
<i>Holarrhena antidysenterica</i> leaves	141±1.50	262±3.3	86.3±1.6	115±4.4	51.7±0.4	31.9±1.1	121±3.51	10.2±0.01
<i>Aegle marmelos</i> leaves	120±2.02	180±4.1	60.4±1.2	100±3.4	41.8±0.7	27.9±1.4	115±4.80	8.21±0.01

<sup>1</sup>Calcium; <sup>2</sup>Phosphorus; <sup>3</sup>Potassium; <sup>4</sup>Magenesium; <sup>5</sup>Zinc; <sup>6</sup>Manganese; <sup>7</sup>Sodium; <sup>8</sup>Copper

Phyto-constituent's in different medicinal plants collected from Orathur village, India (mg/g) is presented in Table 3. Alkaloids, tannins, saponins, flavonoids, terpenoids and phenols in *Tephrosia purpurea*, *Adhatoda vasica*, *Crataeva nurvala*, *Bauhinia variegata*, *Holarrhena antidysenterica* and *Aegle marmelos* leaves ranged from 26.35 – 35.09 mg/g, 39.00 – 46.50 mg/g, 20.05 – 30.40 mg/g, 66.10 – 92.16 mg/g, 40.70 – 45.88 mg/g and 102.1 – 125.4 mg/g respectively. The medicinal property of herbs is due to the presence of different complex chemical substance as secondary metab olites, which are exclusively accumulated in different parts of the plants, (Haniyeh et al., 2010[24]; John, 2024)[31]. Values recorded for tannins, saponins, flavonoids and phenols in this study was lower than values recorded for *Gossypium herbaceum* leaf 69.56 mg/g, 66.67 mg/g, 91.03 mg/g and 219.20 mg/g reported by Oloruntola et al. (2023)[32] but higher than those reported by Falowo et al. (2023)[33] on *Crassocephalun crepidiodes* and *Dysphania ambrosiodes* leaves. This variation could be attributed to age of plant, species, processing method, geographical location amongst others (John, 2024[25]; Ojediran et al., 2024[3]). The result in this experiment suggests that all the plants determined contain several bioactive compounds with therapeutic properties (anti-inflammatory, antioxidant, anti-fungal, antimicrobial, immune-stimulatory, cytotoxic, hepato-protective, anti-viral, anti-helminthic, amongst others) (Agubosi et al., 2022)[13]. For instance, flavonoids and terpenoids have been suggested to possess antioxidant (Yu et al., 2014)[26], antibacterial activity (Edeoga et al., 2005)[27], antimicrobial and anti-cancer properties (Zakariya et al., 2017)[28]. Saponins are reported for hypoglycemic activity, antioxidant activity, cytotoxic and antimicrobial activity (Edeoga et al., 2005)[27]. Olajuyige et al. (2011)[29] reported that alkaloids have analgesics, anti-plasmodial, antiprotozoal and antimicrobial activity. Phenols are reported for antioxidant, anti-cancer, antimicrobial and anti-inflammatory activity (Jing-Chung et al., 2007)[30].

**Table 3** Phyto-constituent's in different medicinal plants collected from Orathur village, India (mg/g)

Medicinal plants	Alkaloids	Tannins	Saponins	Flavonoids	Terpenoids	Phenols
<i>Tephrosia purpurea</i> leaves	26.35±0.61	42.07±1.04	20.05±2.83	86.11±0.05	41.23±0.05	102.1±6.77
<i>Adhatoda vasica</i> leaves	30.91±1.00	40.35±1.88	22.18±2.00	72.03±1.00	40.80±0.07	108.4±5.06
<i>Crataeva nurvala</i> leaves	29.37±1.02	43.11±1.03	30.00±1.83	66.10±0.08	42.19±0.01	113.1±4.11

Bauhinia variegata leaves	35.09±1.10	46.50±1.15	30.40±2.02	92.16±0.02	45.88±0.00	125.4±3.56
Holarrhena antidysenterica leaves	28.55±0.83	41.22±1.97	27.50±1.56	71.90±0.04	40.71±0.02	106.8±3.09
Aegle marmelos leaves	29.63±1.22	39.00±2.31	22.00±1.19	80.11±0.00	43.20±0.01	112.1±1.24

#### 4 CONCLUSION

In conclusion, *Tephrosia purpurea*, *Adhatoda vasica*, *Crataeva nurvala*, *Bauhinia variegata*, *Holarrhena antidysenterica* and *Aegle marmelos* leaves contains several nutritional and medicinal properties at various concentrations which could be attributed to age of plant, species, harvesting method, processing technique amongst other factors. This plants have huge potential in addressing the increasing cases of antimicrobial resistance. It can also be used to promote food safety and livestock production once it is supplemented in their diets.

#### COMPETING INTERESTS

The authors have no relevant financial or non-financial interests to disclose.

#### REFERENCES

- [1] Vera, B. Essential oils Knowledge: from facts to farm or from farm to facts. *International Poultry Magazine*, 2022, (4): 3-4.
- [2] Ojediran TK, Emiola IA, Durojaye V, et al. Proximate, vitamin and GC/MS profiling of *Kigelia africana* fruit powder. *Corrado: Agric. Biotechnology Journal of Science*, 2024, 1(1): 13-20.
- [3] Ojediran TK, Emiola IA, Durojaye V, et al. Analysis of *Kigelia africana* fruits powder antioxidant and phytochemical properties. *Brazilian Journal of Science*, 2024, 3(7): 38-49.
- [4] Inge, H , Rainer, A. Essential oils may reduce the use of antibiotics. *International Poultry Magazine*, 2020, 1(8): 1-2.
- [5] Singh Sharma., Alagbe Olujimi John., et al. Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroemetry. *International Journal of Agriculture and Animal Production*, 2022, 2(6): 18-27.
- [6] John, A.O. Growth performance, haemato-biochemical indices of broiler chicken fed *Aristochia indica* as a phytogetic feed additive. *Cerrado: Agricultural and Biological Research*, 2024, 1(1), 42-53.
- [7] John, A.O., Muritala, S.D., Aduragbemi, Y.A., et al. The approximate mineral and phytochemical content of the leaves, stem bark and roots of *Pterocarpus erinaceus* in India. *Cerrado: Agricultural and Biological Research*, 2024, 1(1), 32-41.
- [8] Musa, B., Alagbe, J.O., Adegbite Motunrade Betty, et al. Growth performance, caeca microbial population and immune response of broiler chicks fed aqueous extract of *Balanites aegyptiaca* and *Alchornea cordifolia* stem bark mixture. *United Journal for Research and Technology*, 2020, 2(2):13-21.
- [9] Ahmad, I., Aqil, F., Ahmad, F, et al. Herbal Medicines: Prospects and Constraints. In: *Modern Phytomedicine: Turning Medicinal Plants into Drugs*, ed. Ahmad I., Aqil F., and Owais M., WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim. 2006, pp. 59-76.
- [10] Aborisade, A.B., Adetutu, A., Owoade, A.O. Phytochemical and proximate analysis of some medicinal leaves. *Clinical Medicine Research*, 2017, 6(6); 209-214.
- [11] Lawal, H.Z., Olatunji, K, Abdulazeez, A. Comparative study on the proximate, mineral and phytochemical analysis of *Daniellia oliveri*, *Leptadenia hastate* and *Vitex doniana* leaves. *Bayero Journal of Pure and Applied Sciences*, 2022, 13(1): 27-33.
- [12] Agubosi, O.C.P., Alexander, James, et al. Influence of dietary inclusion of Sunflower (*Helianthus annus*) oil on growth performance and oxidative status of broiler chicks. *Central Asian Journal of Medical and Natural Sciences*, 2022, 2(7): 187-195.
- [13] Agubosi, O.C.P., Soliu, M.B, Alagbe, J.O. Effect of dietary inclusion levels of *Moringa oleifera* oil on the growth performance and nutrient retention of broiler starter chicks. *Central Asian Journal of Theoretical and Applied Sciences*, 2022, 3(3): 30-39
- [14] Sales, J, Janssens, G. Energy and protein nutrition of companion birds. *Vlaams Diergrneeskundig Tijdschrift*, 2003, 72:51-58.
- [15] NRC. *Nutrient requirements of poultry* (9th Revised Ed.). Washington, DC: National Academy Press, 1994.
- [16] Akaninyene, U.U, Uwemedimo, E.U. Nutritive and anti-nutritive evaluation of *Cnidioscolus aurifolia* leaves. *Turkish Journal of Agriculture – Food Science and Technology*, 2016, 4(9): 739-742.
- [17] Alagbe, J.O. Chemical evaluation of proximate, vitamin and amino acid profile of leaf, stem bark and roots of

- Indigofera tinctoria. International Journal on Integrated Education, 2020, 3(10): 150-157.
- [18] Alagbe, J.O., Adeoye, Adekemi, Oluwatobi, O.A. Proximate and mineral analysis of Delonix regia leaves and roots. International Journal on Integrated Education, 2020, 3(10): 144-149.
- [19] Nahar, M.A., Habiba, U, Sen, P. Evaluation of proximate composition, mineral nutrients, antioxidant and flavonoids and phenolic compounds of Crataeva nurvala leaves and stem bark from Natrokona district of Bangladesh. Journal of Research in Agriculture and Animal Science, 2023, 10(5): 1-6.
- [20] FDA. Nutrition and supplement facts labels: questions and answers related to the compliance date, added sugars, and declaration of quantitative amounts of vitamins and minerals: guidelines for industry retrieved from <https://www.fda.gov/FoodsGuidelines>, 2019.
- [21] Julian, K. What is required daily intake of vitamins and minerals. Medical News Today Magazine, 2021, pp. 23.
- [22] Andzouana, M, Mombouli, J.B. Proximate, mineral and phytochemical analysis of the leaves of Hippocratea myriantha and Urera trinervis. Pakistan Journal of Biological Sciences, 2012, 15(11): 536-541.
- [23] Abolaji, O.A., Adebayo, A.H, Odesanmi, O.S. Nutritional qualities of three medicinal plant parts Xylopia aethiopica, Blighia sapida and Parinari polyandra commonly used by Pregnant women in the Western part of Nigeria. Pakistan Journal of Nutrition, 2007, 6(6): 665-668.
- [24] Haniyeh Koochak, Seyyed Mansour Seyyednejad, Hussein Motamedi. Preliminary study on the antibacterial activity of some medicinal plants of Khuzestan (Iran). Asian Pacific Journal of Tropical Medicine, 2010, 3(3):180-184.
- [25] John, A.O. Clerodendron splendens leaf extract supplementation in weaner rabbits: impact on growth performance, haematology and intestinal microbial population. Cerrado: Agricultural and Biological Research, 2024, 1(1), 21-31.
- [26] Yu EA, Kim GS, Jeong SW, et al. Flavonoid profile and biological activity of Korean citrus varieties (II): Pyunkyul (Citrus tangerina Hort. ex Tanaka) and overall contribution of its flavonoids to antioxidant effect. Journal of Functional Foods, 2014, 6:637-642.
- [27] Edeoga, H.O., Okwu, D.E, Mbaebie, B.O. Phytochemical constituents of some Nigerian medicinal plants. African Journal of Biotechnology, 2005, 4 (7): 685-688.
- [28] Zakariya, R.F., Sani, A., Odebisi-Omokanye, M.B., et al. Screening of Prosopis africana for Antimicrobial Activity Against Selected Oral Pathogens. Nigerian Journal of Pure and Applied Sciences, 2017, 30 (2): 3049-3058.
- [29] Olajuyige, O.O., Babalola, A.E, Afolalayan, A.J. Antibacterial and phytochemical screening of crude ethanolic extracts of Waltheria Linn. African Journal of Microbiology Research, 2011, 5 (22), 3760-3764.
- [30] Jing-Chung, C., Jang-Ying, Y., Pei-Chun, C, et al. Phenolic content and DPPH radical scavenging activity of Yam containing Surimi gels influenced by salt and heating. Asian Journal of Health and Information Sciences, 2007, 2: 1-11.
- [31] John, A.O. Effect of coconut shell extract on the growth performance and some haemato-biochemical parameters of broiler chicken. Brazilian Journal of Science, 2024, 3(6): 82-95.
- [32] Oloruntola, O.D., Simeon, O.A., Victor, A., et al. Evaluation of Gossypium herbaceum leaf powder's nutritional composition and nutraceutical properties. Black Sea Journal of Agriculture, 2023, 7(1): 7-13.
- [33] Falowo, B.F., Oloruntola, O.D, Akinmoladun, O.F. Assessment of nutritional composition and antioxidant properties of Crassocephalun crepidiodes and Dysphania ambrosiodes leaf meal. Turkish Journal of Agriculture-Food Science and Technology, 2023, 11(2): 274-279.
- [34] Alagbe, J.O. Use of medicinal plants as a panacea to poultry production and food security: A review. Gospodarka I Innowacje, 2022, 22(2022): 1-12.
- [35] Musbau, B., Lawal, I.O, Olaniyi, A.A. Proximate and phytochemical screening and mineral analysis of Crescentia cujete leaves. Journal for Medicinal Plants for Economic Development, 2018, 2(1): 1-7.