

PROXIMATE, MINERAL COMPOSITION AND PHYTO-CONSTITUENTS OF SOME MEDICINAL PLANTS/HERBS IN INDIA

Alagbe, JO^{1*}, Anuore, DN², Aliyu, KI²

¹Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India.

²Department of Animal Science, University of Abuja, Nigeria.

Corresponding author: Alagbe, J.O, Email: dralagbe@outlook.com

Abstract: This study was conducted to assess the proximate, mineral content, and phyto-constituents of several therapeutic plants and herbs found in India. The medicinal properties of herbs are attributable to the existence of several complex chemical substances known as secondary metabolites, which are exclusively accumulated in diverse sections of the plant such as leaves, stems, roots, and flowers. These secondary metabolites or phytochemicals contain saponins, alkaloids, flavonoids, triterpenoids, diterpenoids, tannins, and steroids, which are considered a valuable source of nutrition and also possess pharmacological properties such as antimicrobial, antifungal, antiviral, antihelminthic, antioxidant, hepato-protective, antibacterial, immunostimulatory, hypolipidemic, anti-rheumatic, antidiarrheal, anti-pyretic, antimalarial, anticancer, and anti-allergic properties, among others. Medicinal plant extracts have traditionally been used to treat diseases and inhibit the activities of pathogenic organisms such as *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Enterococcus faecalis*, *Staphylococcus aureus*, *Bacillus subtilis*, *Penicillium notatum*, and *Candida albicans*. It was determined that all medicinal plants investigated had varied chemical compositions, which might be ascribed to the plant's age, specie, extraction procedure or processing, storage, geographical location, and other factors. This study will also provide information on emerging phyto-constituents, which can help to lower the growing incidence of antimicrobial resistance and bridge the gap between animal production and food safety.

Keywords: Medicinal plants; Phytochemicals; Toxicity; Treatment; Antimicrobial; Resistance

1 INTRODUCTION

Medicinal or herbal plants have nutritional, pharmacological, or therapeutic characteristics because they contain phytochemicals [1]. They've been used to treat a variety of ailments for thousands of years. The use of terrestrial plants as medicines has been reported in Egypt, China, India and Greece from ancient times, and an astounding number of modern pharmaceuticals have been produced from them [1]. Plants' natural ingredients or phytochemicals can be obtained from their stem barks, leaves, flowers, roots, fruits, and seeds [2]. In 1999, the World Health Organization (WHO) reported that 80% of the world's population used medicinal plants to heal ailments. In recent years, medicinal plants have become a primary health supply for the pharmaceutical business [3].

Medicinal plants are currently gaining acceptance among literates in urban settlements, most likely due to the increasing inefficacy of many modern drugs used to control many infections, as well as the increase in resistance by several bacteria to various antibiotics and the rising cost of prescription drugs for personal health maintenance [4,5]. Current issues with antibiotic use, as well as the rising incidence of multiple-drug resistance strains of a variety of harmful bacteria, have reignited interest in plants having antimicrobial capabilities [6]. Antimicrobial resistance endangers the population by making infections difficult to cure and raising morbidity and mortality rates. It also raises the expense of disease treatment and management [7,8].

Secondary metabolites are the active ingredients in many plant-derived medications. Plant extracts' antibacterial properties may be attributed to a range of components, including aldehydes and phenolic chemicals [9,10]. According to the World Health Organization [11], there are over 100,000 medicinal plant species worldwide, many of which have yet to be identified. They perform multiple biological activities such as: antimicrobial [12,13], antioxidant, anti-inflammatory [14], anti-helminthic, anticancer, antimalarial, anti-rheumatic [11,15], anti-diuretic, anticonvulsant, antidiarrheal, antiviral [16], hepato-protective, immuno-stimulatory, anti-fungal, dermato-protective, hypolipidemic

Infusions of any or extracts from plant parts have also been used traditionally for the treatment of various diseases and infections such as cough, chest, pain, waist pain, irregular menstruation, internal pile, malarial, quick ejaculation, headache, hypertension, dysentery, premature aging, memory improvement, blood cleansing, chronic venous, insufficiency, mental function, minor burns, scars, scleroderma, skin ulcers, varicose veins, wound healing, rheumatism, blood disease, congestive heart failure, urinary tract infections, venereal disease, hepatitis and high blood pressure phlebitis, leg cramps, gastro intestinal diseases, amongst others [17]. They have also been shown to suppress the activity of *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Enterococcus faecalis*, *Staphylococcus aureus*, *Bacillus subtilis*,

Penicillium notatum, and Candida albicans, among others [17,18]. However, factors such as species, age of medicinal plants, and geographical location may impact plant nutritional and phytochemical composition [19]. Given the pharmacological or therapeutic activity associated with various phytochemicals in medicinal plants, it is necessary to identify their chemical components or phyto-constituents, as this could lead to the discovery of novel drugs while also providing information on their harmful effects.

2 MATERIALS AND METHODS

2.1 Experimental Location

The experiment was carried out at Sumitra Research Laboratory Gujarat, India located between 23° 13' N and 72° 41' E. All laboratory kit/machines were operated according to their manufacturer's recommendation.

2.2 The collection, Identification, and Verification of Medicinal Plants

Fresh leaves from 16 medicinal plants were picked from various strands of trees at the Sumitra Research Institute research farm in Gujarat, India. Collected samples were delivered to the institute's Crop Protection Department for proper identification and authentication, with each sample allocated a voucher specimen number. Following that, they were air dried separately for 14 days, ground into powder with an electric blender, and placed in a marked polythene bag for easy identification. The samples were transported to the Sumitra Biological Laboratory for additional analysis.

2.3 Reagents (Analytical Grade) Used for Analysis

Sodium hydroxide, copper sulphate, sulphuric acid, Folin-Ciocalteu reagent, aluminium chloride, sodium carbonate solution, sodium nitrate, isobutyl alcohol, ferric chloride solution, trioxonitrate (v) acid, boric acid, potassium ferricyanide solution, zinc acetate solution, petroleum ether, potassium hydroxide, sodium sulphate, aqueous ammonia solution, chloroform, folin-Denis reagent, hydrochloric acid, Liebermann Buchard reagent, sodium nitroprusside, sodium bicarbonate, bromocresol and pyridine.

2.3 Machines/kit Used for Analysis

Automated Atomic Absorption Spectrometer Skyray (Model; AAS 9000, USA) was used for mineral analysis. GC/MS Ultrospec® 7500, United Kingdom was used for the analysis of phyto-constituents. Automated Near Infra Red Kit (NIR) was used for proximate analysis

2.4 Mineral Analysis of Medicinal Plants

Samples of medicinal plants were analyzed using Skyray instrument (Model: AAS 9000, USA) integrated with atomic absorption spectrometer combining both flame and atomic graphite. To achieve accurate result, 100 g of each sample was passed through the sample thin chamber. Prior to the analysis, 7 lamps in the kit were preheated simultaneously before it was set at a wavelength coverage, repeatability and accuracy of 900nm, $\pm 0.25\text{nm}$ and $<0.10\text{nm}$ respectively and a heating rate of 3000°C/s , static stability of 0.003 Abs according to the manufacturers recommendation before generating results via the visual display unit in less than 120 seconds.

2.5 Gas Chromatography and Mass Spectrophotometer (GC/MS)

Analysis of phyto-constituents in medicinal plants was carried out using Ultrospec® 7500, United Kingdom. Samples was injected through the sample chamber which contains two silicon photodiodes and Xenon flash lamp before it was adjusted at a spectral band width ($<2\text{nm}$), wavelength range (190 – 1100 nm), photometric accuracy and reproducibility of 0.5 % and photometric range of -3.000 – 3.000 A.

2.6 Proximate Analysis of Medicinal Plants

Proximate analysis of medicinal plant was carried out using Phoenix 5000 near infra-red (NIR) analyzer. Kit was calibrated using specific reagents according to the manufacturer's recommendation. 100 g of each sample was placed in a sample cap and the machine was set at a wavelength range, reproducibility and accuracy of 1100 – 2500 nm, $<0.002\text{nm}$ and 0.3 nm, photometric noise ($< 15 \mu\text{Au}$) before results were generated on the monitor in less than 60 seconds.

3 QUANTIFICATION OF PHYTO-CONSTITUENTS IN MEDICINAL PLANTS BY GC/MS

3.1 Total Flavonoid Content Estimation

Total flavonoid content was measured by aluminium chloride colorimetric assay outlined by Tolari *et al.* [20]. 100 grams of each samples were added to 10 ml volumetric flask containing 4 ml of distilled water. To the above mixture, 0.3 ml of 5 percentage sodium nitrate was added followed by aluminium chloride after 30 minutes. Mixture was set in an Ultrospec® 7500 and measured at an optical density of 450 nm.

3.2 Quantification of Total Phenols

The total phenols in the sample was carried out according to methods described by of Makkar *et al.* [21]. 100 grams of each samples were transferred into a test tube followed by the addition of 1 ml of Folin-Ciocalteu reagent and 1 ml of sodium carbonate solution after 30-minute interval before the mixture was set in an Ultrospec® 7500 and measured at an optical density of 550 nm.

3.3 Saponin Estimation

100 g of the grinded dried sample was weighed into a beaker followed by the addition of isobutyl alcohol after 2 minutes. The mixture was shaken and kept in a dark. Thereafter, the mixture was filtered through a filter paper into a beaker containing magnesium carbonate. The solution was mixed well and the absorbance was measured against the prepared reagent blank at 380 nm in an Ultrospec® 7500.

3.4 Total Alkaloid Analysis

100 g of grinded sample was weighed into a beaker containing absolute alcohol. Thereafter, the mixture was transferred to a 100 ml flask followed by the addition of 0.5 g magnesium oxide. The mixture was digested in a boiling water bath for an hour and was filtered while hot through a Buchner funnel. After 30 minutes, 2 drops of alcohol, hydrogen chloride, zinc acetate solution, potassium ferricyanide solution were thoroughly mixed to give a homogenous mixture. Mixture was set in an Ultrospec® 7500 and measured at an optical density of 660 nm.

3.5 Total Tannins Estimation

100 g of sample was measured into a conical flask followed by the addition of methanol and placed in a water bath for 30 minutes. The mixture was shaken. 1.5 mL of Folin-Denis reagent was added with 0.5 mL of sodium bicarbonate before it was filtered into a volumetric flask. The mixture was made up to mark with water mixed well and allowed to stand for 30 minutes. Mixture was set in an Ultrospec® 7500 and measured at an optical density of 350 nm.

3.6 Total Glycosides Analysis

100 g of the sample were measured into 100 mL of conical flask followed by the addition of 5 ml of chloroform, 2 ml of pyridine and sodium nitroprusside shaken thoroughly for 20 minutes before 3 ml of sodium hydroxide was added. Mixture was set in an Ultrospec® 7500 and measured at an optical density of 600 nm.

3.7 Total Steroid Analysis

100 g of each sample were weighed into a conical flask followed by 5ml of chloroform to dissolve the extract and it was kept for 10 minutes. Potassium hydroxide was also added at 2 ml to obtain a homogenous mixture and set in a water bath at 40°C for 60 minutes. Thereafter, 10 ml of petroleum ether and Liebermann Buchard reagent before it was evaporated to dryness. Mixture was set in an Ultrospec® 7500 and measured at an optical density of 720 nm.

3.8 Statistical Analysis

All the tests were carried out in triplicates outcomes were expressed as the mean value \pm standard deviation.

4 RESULTS

Proximate composition of some sixteen medicinal plants found in India (expressed in %) is presented in Table 1. The crude protein values obtained in this study varied from 7.31 – 17.1 %, moisture (8.13 – 11.7 %), crude fat (1.29 – 2.61 %), crude fibre (10.9 – 18.4 %), ash (7.55 – 9.87 %) and carbohydrates (40.9 – 55.6 %). *Centella asiatica* had the highest crude protein value while *Tribulus terrestris* had the lowest value. Higher moisture content was recorded for *Tinospora cordifolia* while *Symplocos racemose* had the lowest value. Result on crude fibre, crude fat, ash and carbohydrate revealed that *Tribulus terrestris*, *Glycyrrhiza glabra*, *Bauhinia variegata* and *Centella asiatica* had a higher value compared to *Crateva nurvala*, *Tribulus terrestris*, *Symplocos racemose* and *Psoralea corylifolia* which had a lower value respectively.

As presented in Table 2, mineral composition of some medicinal plants found in India (expressed in mg/100g). Result revealed that calcium level varied from 155.7 – 211.6 mg/100g, phosphorus (78.9 – 114.2 mg/100g), potassium (398.7 – 831.1 mg/100g), magnesium (47.2 – 96.0 mg/100g), manganese (10.5 – 31.9 mg/100g), zinc (29.1 – 48.0 mg/100g), iron (5.12 – 13.2 mg/100g), selenium (0.43 – 3.11 mg/100g), copper (1.83 – 8.67 mg/100g) and sodium (101 – 132 mg/100g).

As presented in Table 3, phytochemical evaluation of some medicinal plants found in India (expressed in mg/g). Result showed that the concentrations of alkaloids, saponins, flavonoids, tannins, glycosides, steroids and phenols were greater in *Terminalia bellerica* (40.6 mg/g), *Centella asiatica* (45.8 mg/g), *Bauhinia variegata* (70.5 mg/g), *Eclipta alba* (29.5 mg/g), *Andrographis paniculata* (7.12 mg/g), *Bauhinia variegata* (6.17 mg/g) and *Terminalia bellerica* (92.5 mg/g) compared to those of *Psoralea corylifolia* (21.3 mg/g), *Symplocos racemose* (22.0 mg/g), *Withania somnifera* (37.0 mg/g), *Plumbago zeylanica* (16.3 mg/g), *Tinospora cordifolia* (2.06 mg/g), *Psoralea corylifolia* (2.67 mg/g) and *Barleria prionites* (51.4 mg/g) respectively.

Table 1 Proximate Composition of Some Medicinal Plants Found in India (Expressed In %)

Medicinal plants (leaves)	¹ MC	² CP	³ CF	⁴ CFF	⁵ CHO	ASH
<i>Eclipta alba</i>	9.87 ± 0.88	11.4 ± 0.35	16.7 ± 1.46	2.11 ± 0.02	49.8 ± 0.80	7.55 ± 0.19
<i>Andrographis paniculata</i>	10.2 ± 0.65	10.9 ± 0.66	14.5 ± 0.97	1.95 ± 0.01	51.4 ± 0.76	8.04 ± 0.11
<i>Barleria prionites</i>	9.11 ± 0.30	12.6 ± 0.25	16.2 ± 0.88	2.59 ± 0.00	45.6 ± 0.00	8.15 ± 0.04
<i>Withania somnifera</i>	11.4 ± 0.18	14.1 ± 0.17	15.1 ± 0.62	2.03 ± 0.05	42.8 ± 0.11	7.87 ± 0.12
<i>Glycyrrhiza glabra</i>	10.6 ± 0.40	13.6 ± 0.11	16.8 ± 0.85	2.61 ± 0.08	44.0 ± 0.85	9.00 ± 0.06
<i>Tinospora cordifolia</i>	12.8 ± 0.57	15.2 ± 0.70	15.9 ± 0.51	2.55 ± 0.02	53.4 ± 0.73	8.51 ± 0.08
<i>Holarrhena antidysenterica</i>	11.7 ± 0.35	14.3 ± 0.88	17.8 ± 0.70	2.49 ± 0.01	51.2 ± 0.70	8.00 ± 0.00
<i>Plumbago zeylanica</i>	10.6 ± 0.60	16.8 ± 0.40	13.6 ± 0.81	2.00 ± 0.00	49.8 ± 0.69	7.64 ± 0.31
<i>Symplocos racemose</i>	8.13 ± 0.26	7.49 ± 0.10	16.8 ± 1.12	1.96 ± 0.06	46.7 ± 0.85	6.07 ± 0.20
<i>Psoralea corylifolia</i>	10.1 ± 0.11	10.8 ± 0.12	17.2 ± 1.88	1.72 ± 0.04	40.9 ± 0.90	6.42 ± 0.15
<i>Tribulus terrestris</i>	11.0 ± 0.45	7.31 ± 0.06	18.4 ± 0.31	1.50 ± 0.01	55.6 ± 1.00	6.55 ± 0.12
<i>Terminalia arjuna</i>	9.50 ± 0.40	16.2 ± 0.90	16.1 ± 0.82	1.87 ± 0.05	51.2 ± 1.20	6.60 ± 0.05
<i>Crateva nurvala</i>	8.33 ± 0.31	9.04 ± 0.11	10.9 ± 0.46	2.04 ± 0.02	54.0 ± 1.06	8.11 ± 0.12
<i>Terminalia bellerica</i>	9.28 ± 0.12	13.2 ± 0.85	11.5 ± 0.50	1.51 ± 0.04	49.6 ± 0.97	9.26 ± 0.15
<i>Centella asiatica</i>	10.5 ± 0.50	17.0 ± 0.80	12.3 ± 0.11	1.29 ± 0.02	54.7 ± 1.88	9.10 ± 0.04
<i>Bauhinia variegata</i>	10.0 ± 0.61	17.1 ± 0.12	11.6 ± 0.18	1.87 ± 0.12	45.1 ± 1.22	9.87 ± 0.03

¹Moisture content; ²Crude protein; ³Crude fibre; ⁴Crude fat; ⁵Carbohydrates

Table 2 Mineral Composition of Some Medicinal Plants Found in India (Expressed in mg/100g)

Medicinal plants	¹ Ca	² Phos	³ K	⁴ Mg	⁵ Mn	⁶ Zn	⁷ Fe	⁸ Se	⁹ Cu	¹⁰ Na
<i>Eclipta alba</i>	211.6 ± 0.03	108.1 ± 0.10	812.4 ± 0.09	95.3 ± 0.00	15.1 ± 0.11	48.0 ± 0.11	10.4 ± 0.02	0.61 ± 0.001	4.12 ± 0.02	121 ± 1.4
<i>Andrographis paniculata</i>	194.8 ± 0.00	100.5 ± 0.04	706.1 ± 0.06	81.2 ± 0.19	10.5 ± 0.07	44.6 ± 0.10	7.12 ± 0.00	0.43 ± 0.001	3.61 ± 0.01	109 ± 0.81
<i>Barleria prionites</i>	206.8 ± 0.07	114.2 ± 0.16	800.6 ± 0.04	94.1 ± 0.04	17.3 ± 0.12	31.2 ± 0.40	9.33 ± 0.02	0.45 ± 0.000	3.06 ± 0.06	118 ± 4.87
<i>Withania somnifera</i>	188.5 ± 0.21	106.3 ± 0.10	658.2 ± 0.23	82.6 ± 0.00	21.0 ± 0.90	29.5 ± 0.10	8.56 ± 0.05	0.87 ± 0.002	3.11 ± 0.05	105 ± 3.11
<i>Glycyrrhiza glabra</i>	203.1 ± 0.23	100.2 ± 0.06	811.5 ± 0.15	96.0 ± 0.03	26.4 ± 0.19	30.6 ± 0.04	10.4 ± 0.07	0.68 ± 0.003	2.41 ± 0.03	109 ± 2.86
<i>Tinospora cordifolia</i>	174.9 ± 0.12	98.5 ± 0.12	604.7 ± 0.00	71.0 ± 0.16	23.9 ± 0.06	29.1 ± 0.00	8.55 ± 0.02	0.59 ± 0.006	2.06 ± 0.06	114 ± 3.11
<i>Holarrhena antidysenterica</i>	185.4 ± 0.07	75.9 ± 0.01	516.2 ± 0.90	69.4 ± 0.24	16.5 ± 0.02	31.4 ± 0.56	9.00 ± 0.01	0.74 ± 0.005	2.33 ± 0.05	112 ± 2.00
<i>Plumbago zeylanica</i>	200.1 ± 0.00	93.6 ± 0.00	831.1 ± 0.06	94.2 ± 0.16	20.9 ± 0.01	33.8 ± 0.09	8.60 ± 0.03	0.69 ± 0.004	2.60 ± 0.01	116 ± 3.26

<i>Symplocos racemose</i>	161.0 ±0.01	88.7±0.09	502.8±0.53	56.9±0.04	24.7±0.02	30.7±0.51	6.06±0.06	0.57±0.001	1.83±0.02	101±1.69
<i>Psoralea corylifolia</i>	196.1 ±0.14	90.4±0.15	489.3±0.06	52.1±0.16	12.9±0.00	27.6±0.05	5.67±0.08	0.82±0.000	2.94±0.09	115±1.22
<i>Tribulus terrestris</i>	172.5 ±0.10	81.6±0.40	420.7±0.18	48.4±0.03	18.7±0.09	31.2±0.09	5.12±1.00	0.67±0.009	3.71±0.01	104±1.34
<i>Terminalia arjuna</i>	182.9 ±0.08	78.9 ±0.43	400.6±0.04	50.6±0.31	16.5±0.04	30.4±0.01	6.08±1.22	0.93±0.006	3.03±0.03	112±1.63
<i>Crateva nurvala</i>	155.7 ±0.11	70.2±0.18	398.7±0.12	47.2±0.47	18.4±0.31	32.0±0.07	6.77±1.50	0.67±0.001	5.48±0.00	105±1.24
<i>Terminalia bellerica</i>	194.8 ±0.12	84.1±0.43	408.5±0.08	60.8±0.21	31.2±0.12	38.4±0.12	12.6±1.88	3.11±0.007	7.91±0.00	132±1.34
<i>Centella asiatica</i>	205.1±0.03	93.5±0.12	800.6±0.12	78.9±0.23	30.9±0.18	39.8±0.03	13.8±1.24	2.51±0.005	8.06±0.06	130±2.98
<i>Bauhinia variegata</i>	211.4 ±0.12	95.7±0.00	815.0±0.00	82.6±0.09	31.9±0.04	36.5±0.00	13.2±1.45	2.33±0.001	8.67±0.05	131±2.55

¹Calcium; ²Phosphorus; ³Potassium; ⁴Magnesium; ⁵Manganese; ⁶Zinc; ⁷Iron; ⁸Selenium; ⁹Coppe

Table 3 Phytochemical Evaluation of Some Medicinal Plants Found in India (expressed in mg/g)

Medicinal plants (leaves)	Alkaloids	Saponins	Flavonoids	Tannins	Glycosides	Steroids	Phenols
<i>Eclipta alba</i>	27.1±0.09	34.0±0.01	51.8±2.06	29.5±1.22	5.40±0.03	5.66±0.00	56.4±1.45
<i>Andrographis paniculata</i>	25.4±0.06	35.1±0.00	45.6±1.21	25.0±0.33	7.12±0.00	4.09±0.01	53.0±1.00
<i>Barleria prionites</i>	23.1±0.01	37.5±0.07	40.9±1.34	26.2±1.43	5.77±0.01	6.60±0.02	51.4±1.13
<i>Withania somnifera</i>	21.9±0.05	38.9±0.06	37.0±1.61	24.9±1.00	6.06±0.03	2.56±0.03	50.9±1.31
<i>Glycyrrhiza glabra</i>	20.6±0.03	31.5±0.02	45.7±0.95	22.0±0.42	4.42±0.05	3.07±0.01	55.6±1.09
<i>Tinospora cordifolia</i>	25.1±0.01	33.8±0.08	48.5±1.12	22.3±0.21	2.06±0.01	4.12±0.02	65.7±1.46
<i>Holarrhena antidysenterica</i>	23.6±0.05	34.0±0.11	42.3±0.45	23.7±0.53	4.11±0.00	5.06±0.00	62.1±1.00
<i>Plumbago zeylanica</i>	26.7±0.00	31.3±0.17	47.6±1.21	16.3±0.41	5.67±0.03	6.31±0.01	60.9±1.70
<i>Symplocos racemose</i>	26.8±0.02	22.0±0.06	47.0±0.46	26.2±0.34	6.07±0.02	6.09±0.03	56.0±2.56
<i>Psoralea corylifolia</i>	21.3±0.21	31.6±0.02	45.1±0.33	21.0±0.25	4.61±0.01	2.67±0.02	55.1±2.21
<i>Tribulus terrestris</i>	22.0±0.12	31.0±0.08	43.8±1.35	21.2±0.00	5.78±0.18	5.31±0.01	56.7±1.21
<i>Terminalia arjuna</i>	24.6±0.01	31.0±0.01	45.6±1.41	24.5±0.16	6.90±0.80	5.00±0.01	59.4±1.00
<i>Crateva nurvala</i>	21.5±0.83	32.7±0.03	47.9±1.04	22.3±0.18	4.33±0.93	5.32±0.03	67.3±0.09
<i>Terminalia bellerica</i>	40.6±0.07	42.0±0.04	67.1±0.06	27.0±0.33	3.94±0.90	5.08±0.01	92.5±0.00
<i>Centella asiatica</i>	38.7±0.01	45.8±0.31	65.6±0.04	26.2±0.12	3.40±0.16	6.00±0.03	90.6±0.02
<i>Bauhinia variegata</i>	32.8±0.04	40.9±0.32	70.5±0.51	28.3±0.34	4.52±0.70	6.17±0.01	91.5±0.15

5 DISCUSSION

According to Alagbe (2019a) Alagbe *et al.* (2019b)[22,23], a plant's chemical makeup can be influenced by factors such as age, geographical location, and species. The crude protein range (7.31 - 17.1%) in this study was consistent with Gonzalez *et al.*'s established values for leafy vegetables (8.00 - 30.0%)[24]. The findings imply that these therapeutic plants provide both nutritional and pharmacological benefits, making them suitable for inclusion in animal diets. Andrew *et al.* reported protein concentrations for *Dysphania ambrosoides* (15.99%) and *Crassocephalum crepidioides* leaves (17.11%) that are consistent with the investigation[25]. Alagbe *et al.* found a lower value of 7.60% for *Pterocarpus erinaceus* leaves, 5.11% for *Pterocarpus erinaceus* stem bark, and 4.33% for the root[26]. Mohammad *et al.* recorded a higher crude protein of 28.20 % for *Vernonia amygdalina* leaves[27]. Protein are needed for the growth and maintenance of animals, they also play an integral part in the immune system and enzyme production in the body [28]. The range of moisture content 8.13 – 11.7 % was within the standard range of 6 – 15 % reported for vegetables [29]. Therefore, high moisture content in the samples are advantageous because inhibits the growth of microbes on a sample [30]. The difference in moisture content between different plants is directly dependent on climatic changes [24,60]. This result corresponds to the reported values by Adewale *et al.* [30] for *Pterocarpus carpus* leaves (11.46 %) and stem bark (13.18 %) and lower when compare with the report of Alagbe *et al.* (2022) for *Piliostigma thonningii* leaves (7.23 %). The results on the crude fibre suggests that *Tribulus terrestris* had the highest concentration (18.4 %). Availability of such high contents in the diets of livestock's helps to

prevent gastrointestinal disorders and coronary heart disease [31,32]. The crude fibre range (10.4 – 18.9 %) were within the range reported by Abiodun et al. [33] for *Chenopodium ambrosoides* leaves (13.40 %) and *Morinda lucida* leaves (15.06 %) but lower than values recorded for *Parquetin nigrescens* leaves (22.05 %), *Oscium gratissimum* (22.02 %) and *Magnifera indica* (19.01 %) by the same author. Ash content of medicinal plants can be used to evaluate its mineral content [34,35]. The ash content range reported in this study 7.55 – 9.87 % showed that *Bauhinia variegata* had the highest mineral content suggesting that it can promote construction of muscles, blood cells, internal organs and enzymes in animals [36]. Therefore, animals with a deficiency of minerals will never develop properly and are more susceptible to diseases [37]. The mineral content of medicinal plants is in consonance with the reports of Amabye [38] for *Rhamnus prinoides* leaves (9.50 %) and lower when compared with the reports of Raimi et al. for *Sida acuta* leaves (6.33 %)[39]. Crude fat content range of 1.29 – 2.61 % recorded for the medicinal plants indicated that *Terminalia arjuna* had the lowest value. This result suggests that they are able to maintain good health by preventing the incidence of cardiovascular diseases as a result of excessive fat consumption [40]. Carbohydrate values (40.9 – 55.6 %) corresponds with the values of Abiodun et al. [33] on *Oscium gratissimum* (50.06 %) and lower when compared with the report of the same author for *Parquetin nigrescens* leaves (36.03 %), *Chenopodium ambrosoides* (4.36 %) and *Magnifera indica* (43.76 %). Availability of high carbohydrate content in a sample is helpful in providing energy and excess of it can be stored as fat which is stored in the adipose tissues of animals.

Calcium, phosphorus, potassium, magnesium, manganese, zinc, iron, selenium, copper and manganese in *Eclipta alba* (211.6 mg/100g), *Barleria prionites* (114.2 mg/100g), *Plumbago zeylanica* (831.1 mg/100g), *Glycyrrhiza glabra* (96.0 mg/100g), *Terminalia bellerica* (31.2 mg/100g), *Terminalia bellerica* (3.11 mg/100g), *Bauhinia variegata* (8.67 mg/100g) and *Terminalia bellerica* (132.0 mg/100g) compared to those of *Crateva nurvala* (155.7 mg/100g), *Crateva nurvala* (70.2 mg/100g), *Crateva nurvala* (398.7 mg/100g), *Crateva nurvala* (47.2 mg/100g), *Andrographis paniculata* (10.5 mg/100g), *Psoralea corylifolia* (27.6 mg/100g), *Tribulus terrestris* (5.12 mg/100g), *Andrographis paniculata* (0.43 mg/100g), *Symplocos racemose* (1.83 mg/100g) and *Symplocos racemose* (101 mg/100g) which had lower values. The variation in their compositions may be due to differences in geographical location and species of plants [41]. The result obtained showed that the medicinal plants contains appreciable quantities of minerals which are needed for the activation of enzymatic reaction in the body of animals [28]. Deficiency of minerals are known to affect the performance and health in both humans and animals [42]. Calcium and phosphorus are essential components of the skeleton and are necessary for the synthesis of structural proteins [41]. The calcium and phosphorus levels are within the standard range reported by Obazelu et al. [43] for *Combretum platypterum* leaves (266.4 mg/100g) but lower than values reported for *Ficus capensis* leaves 186.0 mg/100g and 160 mg/100 respectively by Ngozi et al. [44]. Magnesium activates enzyme systems that maintain electrical potential in nerves, whereas potassium influences osmotic pressure and contributes to normal acid-base balance [42]. Iron is essential in mammalian diet to prevent anemia, and it is a component of hemoglobin and myoglobin molecules that transport oxygen to and within cells. Zinc creates metalloproteinase and enzyme complexes that cannot be separated without losing activity [19,45]. They also help cells replicate and differentiate [46].

Phytochemicals are basically divided into two groups that are primary and secondary metabolites. Primary metabolites comprise common sugar, amino acids, proteins and chlorophyll while secondary metabolites consist of alkaloids, flavonoids, tannins etc., [47]. Phytochemicals like flavonoids and phenols are powerful antioxidants and have an important role in the health care system [28]. The values of alkaloids, saponins, flavonoids, tannins, glycosides, steroids and tannins which varied from 20.6 – 40.6 mg/g, 22.0 – 42.0 mg/g, 37.0 – 70.5 mg/g, 16.3 – 29.5 mg/g, 2.06 – 7.12 mg/g, 2.56 – 6.60 mg/g and 50.9 – 92.5 mg/g respectively. Alkaloid, saponin, tannin and flavonoid concentrations were within the range reported by Aliyu et al. for the leaves of *Anchomanes difformis* (23.6 mg/g)[48], *Anisopus mannii* (41.6 mg/g), *Pavetta crassipes* (39.6 mg/g), *Stachytarpheta augustifolia* (40.8 mg/g) and *Vernonia blumeoides* (38.2 mg/g) but lower than values recorded by Obazelu et al. for *Combretum platypterum* (3.52 mg/g)[43]. The role of medicinal plants in disease prevention or control has been attributed to antioxidant properties of their active constituents [49,50]. Medicinal plants have been shown to function as free radical scavengers through anti-oxidative mechanisms mediated by polyphenols, flavonoids, ascorbic acid, and terpenoids, which have the ability to protect cell organelles from damage caused by free radical-induced oxidative stress by inhibiting the initiation or propagation of oxidative chain reactions [51]. Alkaloids have been shown to act as painkillers as well as having antimalarial and antibacterial activities [52,53]. *Melaleuca Alternifolia* [54], *Cassia Occidentalis* [55], *Rhamnus Californica*, and *Umbellularia Californica* leaves have all been found to contain alkaloids and saponins [56]. Tannins are employed as antiseptics due to the presence of phenolic groups, and they have also been linked to antibacterial and antioxidant characteristics. Saponins are also useful therapeutically because they have been proven to have hypolipidemic and anticancer properties [56]. Glycosides have a bitter taste and have been employed as flavoring ingredients in numerous pharmacological products [57,58]. These natural metabolites are significant as prospective antibacterial crude drugs and a source of natural chemicals as new anti-infection agents [59].

6 CONCLUSION

Medicinal plants are thought to be safe and free of adverse effects when used to improve animal performance. Herbs can be used as feed additives because they are suitable and preferred, have a minimal risk of toxicity, pose few health risks, are

environmentally friendly, and are less expensive to produce. Differences in processing processes, species, geographical regions, plant age, and storage, among other factors, can all have an impact on the chemical composition of medicinal plants. Phytochemicals are natural substances that are easily digested by animals. They can also assist to slow the rise in antimicrobial resistance and improve food safety. They can boost feed intake and digestibility in animals while causing no negative effects or withdrawal periods, enhancing farmer profits.

COMPETING INTERESTS

The authors have no relevant financial or non-financial interests to disclose.

REFERENCES

- [1] Singh N, Singh JP, Singh V. Effect of dietary supplementation of herbal formulation on dressing percentage and mortality in broiler chicks. *Indian J. Vet.*, 2002, 2: 22-24.
- [2] Wojcikowski, K, Johnson, DW, Gobé, G. Medicinal herbal extracts -- renal friend or foe? Part one: the toxicities of medicinal herbs. *Nephrology*, 2004, 9(5): 313-318.
- [3] Phillipson, JD. Assays for antimalarial and amoebicidal activities. In: *Methods in plant biochemistry*, ed. Hostettmann K., Academic Press Limited, Great Yarmouth. Norfolk, 1991, 135-152.
- [4] Smolinski, MS, Hamburg, MA, Lederberg, J. (eds). *Microbial threats to health: Emergence, detection, and response*. Washington, DC: Institute of Medicine, National Academies Press, 2003, 203-210.
- [5] Tzortzakis, NG, Economakis, CD. Antifungal activity of lemongrass (*Cymbopogon citratus* L.) essential oil against key postharvest pathogens. *Innovative Food Science Emerging Technology*, 2007, 8: 253-258.
- [6] Voravuthikunchai, SP, Kitpipit, L. Activities of crude extracts of Thai medicinal plants on methicillin-resistant *Staphylococcus aureus*. *Journal of Clinical Microbiology and Infection*, 2003, 9: 236.
- [7] Zhang, Y, Liu, X, Wang, Y, et al. Antibacterial Activity and Mechanism of Cinnamon Essential Oil Against *Escherichia Coli* and *Staphylococcus Aureus*. *Food Control*, 2016, 59, 282-289. DOI: 10.1016/j.foodcont.2015.05.032.
- [8] Winkel-Shirley, B. Update on flavonoid biosynthesis: Flavonoid Biosynthesis. A Colorful Model for Genetics, Biochemistry, Cell Biology, and Biotechnology. *Plant Physiology*, 2001, 126(2): 485- 493.
- [9] Yadav, J, Arya, V, Yadav, S, et al. *Cassia Occidentalis* L.: A Review on its Ethnobotany, Phytochemical and Pharmacological Profile. *Fitoterapia*, 2010, 81(4): 223-230.
- [10] Valgas, C, Souza, SMD, Smânia, EF, et al. Screening Methods to Determine Antibacterial Activity of Natural Products. *Brazilian Journal of Microbiology*, 2007, 38(2): 369-380.
- [11] World Health Organization, (WHO). *Promoting the Role of Traditional Medicine in Health Systems: A Strategy for the African Region 2001-2010*. Harare, World Health Organization, 2000, 10- 286.
- [12] Daniel Nnadozie Anorue, Friday Ubong, Alagbe Olujimi John. Investigating the effects of pawpaw (*Carica papaya*) essential oil dietary supplementation on the growth performance and carcass characteristics of broilers. *Agricultural and Veterinary Sciences*, 2023, 7(3): 164-174.
- [13] Musa, B, Alagbe, JO, Adegbite Motunrade Betty, et al. Growth performance, caeca microbial population and immune response of broiler chicks fed aqueous extract of *Balanites aegyptiaca* and *Alchornea cordifolia* stem bark mixture. *United Journal for Research and Technology*, 2020, 2(2):13-21.
- [14] Demain, AL, Fang, A. *The Natural Functions of Secondary Metabolites. History of Modern Biotechnology I*. ed. by Anon: Springer, 2000, 69, 1-39. DOI: https://doi.org/10.1007/3-540-44964-7_1.
- [15] Daglia, M. Polyphenols as Antimicrobial Agents'. *Current Opinion in Biotechnology*, 2012, 23(2): 174-181.
- [16] Ayanwuyi, LO, Yaro, AH, Abodunde, OM. Analgesic and Anti-Inflammatory Effects of the Methanol Stem Bark Extract of *Prosopis Africana*. *Pharmaceutical Biology*, 2010, 48(3): 296-299.
- [17] Trentin, DD, Giordani, RB, Zimmer, KR, et al. Potential of Medicinal Plants from the Brazilian Semi-Arid Region (Caatinga) Against *Staphylococcus Epidermidis* Planktonic and Biofilm Lifestyles. *Journal of Ethnopharmacology*, 2011, 137(1): 327-335.
- [18] Yasunaka, K, Abe, F, Nagayama, A, et al. Antibacterial Activity of Crude Extracts from Mexican Medicinal Plants and Purified Coumarins and Xanthones. *Journal of Ethnopharmacology*, 2005, 97(2): 293-299.
- [19] Alagbe, JO, Shittu, MD, Ushie, FT. GC-MS analysis of methanolic stem bark extract of *Zollingeriana indigofera*. *Asian Journal of Advances in Research*, 2021, 11(4): 144-146.
- [20] Tolari S, Rudroju S, Penchala S, et al. Quantification of total phenolics and Total Flavonoid contents in extracts of *Oroxylum Indicum* I. *Kurz. Asain Journal of Pharmaceutical and Clinical Research*, 2012, 5(4): 177-179.
- [21] Makkar HPS, Norsambu T, Lkhavatsere, S, et al. Plant Secondary metabolites in some medical plants of Mongolia used for enhancing animal health and production. *Tropicultura*, 2009, 27 (3): 159-167.
- [22] Alagbe, JO. Growth performance and haemato-biochemical parameters of broilers fed different levels of *Parkia biglobosa* leaf extracts. *Academic Journal of Life Sciences*, 2019, 5(12): 107-115.

- [23] Alagbe, JO, Sadiq, MR, Anaso, EU, et al. Efficacy of Albizia lebbeck seed oil on the growth performance and carcass characteristics of weaner rabbits. *Sumerianz Journal of Agriculture and Veterinary*, 2019, 2(12): 116-122.
- [24] González-Lamothe, R, Mitchell, G, Gattuso, M, et al. (2009) 'Plant Antimicrobial Agents and their Effects on Plant and Human Pathogens'. *International Journal of Molecular Sciences*, 2009, 10(8): 3400-3419.
- [25] Andrew, F, Olugbenga, OD, Oluwakamisi, FA. Assessment of Nutritional and Antioxidant properties of *Dysphania ambrosioides* and *Crassocephalum crepidioides* leaf meal as potential feed additives. *Turkish Journal of Agriculture Food Science and Technology*, 2022, 11(2): 274-279.
- [26] Alagbe, JO, Muritala, SD, Aduragbemi, YA, et al. The approximate mineral and phytochemical content of the leaves, stem bark and roots of *Pterocarpus erinaceus* in India. *Cerrado: Agricultural and Biological Research*, 2024, 1(1): 32-41.
- [27] Mohammad, A, Lurwan, M, Sani, U, et al. Determination of proximate, phytochemicals and mineral composition of *Verononia amygdalina* leaves. *Nutraceutical Research*, 2020, 1(1): 1-7.
- [28] Ojediran, TK, Emiola, IA, Durojaye, V, et al. Proximate, vitamin and GC/MS profiling of *Kigelaia africana* fruit powder. *Cerrado: Agricultural and Biological Research*, 2024, 1(1): 13-20.
- [29] Alagbe Olujimi John. Novel phytochemicals' impact on weaned pigs growth performance, haematology and serum biochemical indicators. *Black Sea Journal of Agriculture*, 2024, 7(2): 82-89.
- [30] Adewale, EF, Oluremi, A, Saheed, OA. Nutrients, phytochemical, antioxidants and antimicrobial analysis of *Pterocarpus* stem bark and leaf for their nutritional and medicinal capacity. *Indonesian Journal of Chemical Research*, 2022, 10(1): 58-67.
- [31] Lagnika, Anago, E, Sanni, A. Screening for Antibacterial, Antioxidant Activity and Toxicity of some Medicinal Plants used in Benin Folkloric Medicine'. *Journal of Medicinal Plants Research*, 2011, 5(5): 773-777.
- [32] Dai, J, Mumper, RJ. Plant Phenolics: Extraction, Analysis and their Antioxidant and Anticancer Properties. *Molecules*, 2010, 15(10): 7313-7352.
- [33] Abiodun, BA, Adewale, A, Abiodun, AA. Phytochemical and proximate analysis of some medicinal leaves. *Clinical Medicine Research*, 2017, 6(6): 209-214.
- [34] Daniel Nnadozie Anuore, Shittu Muritala Daniel, Emiola Adewale, et al. Impact of dietary supplementation of *Carica papaya* essential oil on the blood chemistry of broiler chickens. *Science Letters*, 2023, 11(3): 105-110.
- [35] Alagbe, JO. Use of medicinal plants as a panacea to poultry production and food security: A review. *Gospodarka I Innowacje*, 2022, 22: 1-12.
- [36] Singh Sharma, Alagbe Olujimi John, Liu Xing, et al. Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroemetry. *International Journal of Agriculture and Animal Production*, 2022, 2(6): 18-27.
- [37] Alagbe, JO. *Clerodendron splendens* leaf extract supplementation in weaner rabbits: impact on growth performance, haematology and intestinal microbial population. *Cerrado: Agricultural and Biological Research*, 2024, 1(1): 21-31.
- [38] Amabye TG. Evaluation of Phytochemical, Chemical Composition, Antioxidant and Antimicrobial Screening Parameters of *Rhamnus prinoides* (Gesho) Available in the Market of Mekelle, Tigray, Ethiopia. *Nat Prod Chem Res*, 2015, 4: 198. DOI:10.4172/2329-6836.1000198.
- [39] Raimi, MM, Oyekanmi, AM, Adegoke, BM. Proximate, phytochemical and micro-nutrient composition of *Sida acuta*. *IOSR Journal of Applied Chemistry*, 2014, 7(2): 93-98.
- [40] Mboto, C, Eja, M, Adegoke, A, et al. Phytochemical Properties and Antimicrobial Activities of Combined Effect of Extracts of the Leaves of *Garcinia Kola*, *Vernonia Amygdalina* and Honey on some Medically Important Microorganisms. *African Journal of Microbiology Research*, 2009, 3(9): 557-559.
- [41] Alagbe, JO. Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phytogenic Feed Additive in Growing Rabbit Diets: Growth Performance and Caecal Microbial Population. *Concept in Dairy and Veterinary Sciences*, 2021, 4(2).
- [42] Thomas, RA, Krishnakumari S. Proximate analysis and mineral composition of *Myristica fragrans* seeds *Journal of Pharmacognosy and Phytochemistry*, 2015, 3(6): 39-42.
- [43] Obazelu, PA, Aruomaren, A, Ugboaja, EE. Phytochemical analysis nutrients and mineral composition of *Combretum platypterum* aqueous extract. *Journal of Applied Science and Environment Management*, 2021, 25(9): 1625-1630.
- [44] Ngozi Kalu A, Chimaraoke, O, Chima, AE, et al. Phytochemical, Proximate Analysis, Vitamin and Mineral Composition of Aqueous Extract of *Ficus capensis* leaves in South Eastern Nigeria. *Journal of Applied Pharmaceutical Science*, 2017, 7(3): 117-122.
- [45] Adewale, AO, Alagbe, JO, Adeoye, Adekemi. O. Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phytogenic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. *International Journal of Orange Technologies*, 2021, 3(3): 1-12.
- [46] Robert, KM, Daryl, KG, Peter, AM, et al. Harper's Illustrated Biochemistry. In Benders and Mayes Vitamins and Minerals, Lange Medical Books/McGraw-Hill, Medical Publishing Division, New York, 2003, 496.
- [47] Tzakou, O, Pitarokili, D, Chinou, IB, et al. Composition and antimicrobial activity of essential oil of *Salvia ringens*. *Planta Medica*, 2001, 67: 81-83.

- [48] Aliyu, A, Musa, AM, Oshanmi, JA, et al. Phytochemical analysis and mineral elements composition of some medicinal plants of Northern Nigeria. *Nigerian Journal of Pharmaceutical Science*, 2008, 7(1): 119-125.
- [49] Alagbe, JO, Adeoye, Adekemi, Oluwatobi, OA. Proximate and mineral analysis of *Delonix regia* leaves and roots. *International Journal on Integrated Education*, 2020, 3(10): 144-149.
- [50] Muritala, Daniel Shittu, Alagbe, JO, Ojebiyi, OO, et al. Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of *Polyalthia longifolia* leaf extract in place of conventional antibiotics. *Animal Science and Genetics*, 2022, 18(2): 57-71.
- [51] Shittu, MD, Alagbe, JO. Phyto-nutritional profiles of broom weed (*Sida acuta*) leaf extract. *International Journal of Integrated Education*, 2020, 3(11): 119-124.
- [52] Alagbe, JO. Proximate, mineral and phytochemical analysis of some medicinal plants collected from Orathur village, Thiruporur Taluk Kancheepuram district Tamilnadu, India. *World Journal of Agriculture and Forestry Sciences*, 2024, 2(2): 30-35.
- [53] Alagbe, JO, Sharma, R, Eunice Abidemi Ojo, et al. Chemical evaluation of the proximate, minerals, vitamins and phytochemical analysis of *Daniellia oliveri* stem bark. *International Journal of Biological, Physical and Chemical Studies*, 2020, 2(1):16-22.
- [54] Cox, S, Mann, C, Markham, J, et al. The Mode of Antimicrobial Action of the Essential Oil of *Melaleuca Alternifolia* (Tea Tree Oil). *Journal of Applied Microbiology*, 2000, 88(1): 170-175.
- [55] Chukwujekwu, J, Coombes, P, Mulholland, D, et al. Emodin, an Antibacterial Anthraquinone from the Roots of *Cassia Occidentalis*'. *South African Journal of Botany*, 2006, 72(2): 295-297.
- [56] Carranza, MG, Sevigny, MB, Banerjee, D, et al. Antibacterial Activity of Native California Medicinal Plant Extracts Isolated from *Rhamnus Californica* and *Umbellularia Californica*'. *Annals of Clinical Microbiology and Antimicrobials*, 2015, 14(1): 29-33.
- [57] Alagbe, JO, Shittu, MD, Eunice Abidemi Ojo. Prospect of leaf extracts on the performance and blood profile of monogastric – A review. *International Journal of Integrated Education*, 2020, 3(7): 122-127.
- [58] Madhu M, Sailaja V, Satyadev T NVSS, et al. Quantitative phytochemical analysis of selected medicinal plant species by using various organic solvent. *Journal of Pharmacognosy Phytochemistry*, 2016, 5(2): 25-29.
- [59] Shittu, MD, Alagbe, JO, Adejumo, DO, et al. Productive Performance, Caeca Microbial Population and Immune-Modulatory Activity of Broiler Chicks Fed Different Levels *Sida Acuta* Leaf Extract in Replacement of Antibiotics. *Bioinformatics and Proteomics Open Access Journal*, 221, 5(1): 000143.
- [60] Inngjerdingen, K, Nergård, CS, Diallo, D, et al. An Ethnopharmacological Survey of Plants used for Wound Healing in Dogonland, Mali, West Africa. *Journal of Ethnopharmacology*, 2004, 92(2-3): 233-244.